

MICROCOPY

CHART

COHERENT SCATTERING OF LIGHT BY CRYSTALS

Final Report

to

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH

J. Weber

University of Maryland, College Park, Maryland

and

University of California, Irvine, California

1985





Approved for public released & Distribution Unlimited

UNIVERSITY OF MARYLAND
DEPARTMENT OF PHYSICS AND ASTRONOMY
COLLEGE PARK, MARYLAND

	REPORT DOCUM	NTATION PAGE	E		
1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS			
28. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT			
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE		Approved for public release; distribution unlimited.			
4. PERFORMING ORGANIZATION REPORT NUMBER(S)		5. MONITORING ORGANIZATION REPORT NUMBER(S)			
83NP301	AFOSR-TR. 86-0239				
Dept. of Physics & Astronomy	бь. OFFICE SYMBOL (If applicable)	78. NAME OF MONITORING ORGANIZATION			
Gc. ADDRESS (City, State and ZIP Code) University of Maryland College Park, Maryland 20742		7b. ADDRESS (City,	State and ZIP Cod	<i>r7_0</i> e)	
8. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR/NP	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR-82-0164			
8c. ADDRESS (City, State and ZIP Code)		10. SOURCE OF FU			
Building 410 Bolling AFB, D.C. 20332		PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT NO.
11. TITLE (Include Security Classification) COHERENT SCATTERING OF LIGHT BY CRYSTALS		61102F	2301	A8	
12. PERSONAL AUTHORIS) J. Weber					
Final Scientific FROM 2/1/84 to 1/31/		14. DATE OF REPORT (Yr., Mo., Day) Due 30 March 1985 3			
16. SUPPLEMENTARY NOTATION			-		
17. COSATI CODES			A : A . A		
17. COSATI CODES FIELD GROUP SUB. GR.	18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)				
19. ABSTRACT (Continue on reverse if necessary and	identify by block numbe	r)			
Experiments have bee from a helium neon laser Collective absorptio	interacts with	nuclear spins	in a cryst	al.	
are polarized by a magnet	ic field, and a	ppropriate po	larized lig	ht employed	e spins
20. DISTRIBUTION/AVAILABILITY OF ABSTRACTURE LASSIFIED/UNLIMITED TENTO SAME AS RPT.		21. ABSTRACT	CLASSIFIE	CATION	
220. NAME OF RESPONSIBLE INDIVIDUAL		22b. TELEPHONE N		22c. OFFICE SYM	80L

COHERENT SCATTERING OF LIGHT BY CRYSTALS

Final Report

CONTROL BURBLACE WORKS RESERVE

to

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH

J. Weber

University of Maryland, College Park, Maryland

and

University of California, Irvine, California

1985

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (APSC NOTICE OF TRANSMITTAL TO DEID

approved:

Oistrib.

AATTHEW J. Kanera
Chief, Technical Information Division

ABSTRACT

Experiments have been carried out at liquid helium temperatures, in which light from a helium neon laser interacts with nuclear spins in a crystal.

Collective absorption of photons by the spin system is observed, if the spins are polarized by a magnetic field, and appropriate polarized light employed.

	Accession For	_			
	NTIS GRAMI				
	DTIC TAB				
	Uneppercord [1]				
	Just Mark Comment				
	The second secon				
	Ву				
*	District Continues.				
J. 25	Availability Comes				
`	The state of the s				
	Dist special				
	U·I				

INTRODUCTION

For the past several years we have been studying coherent scattering of red light from a Helium Neon Laser, by sapphire and lithium fluoride crystals. The theory has been presented in earlier proposals, and reports. This theory predicts that nearly perfect crystals with nuclei having magnetic moments, will absorb light by a collective spin state change. Thus many nuclei change their spin state in a magnetic field, absorbing a single red photon.

Experiments

Figure 1 shows the experimental arrangement. Light from the Laser interacts with a crystal in a magnetic field at low temperatures. A long period, about 36 hours, is required for the nuclear spins to achieve thermal equilibrium with the lattice. It is therefore expected that the crystal will be transparent initially and Lecome opaque on a time scale of the hours required for the spin polarization to approach equilibrium values.

Experiments During the Current Grant

Theory predicts that the light must be polarized with magnetic field vector parallel to the applied field which polarizes the nuclei. Early experiments employed an unpolarized laser together with a polarizer, and these gave consistent results for an extended period.

During the present grant period this unpolarized laser was again employed. Data were erratic and not reproducible. A series of experiments revealed that an "unpolarized" laser does age and its output may be polarized with an uncertain variable direction.




FIGURE 1

The laser was then replaced by a polarized one milliwatt laser.

Two neutral density attenuator filters and a diffuser were employed to illuminate the entire crystal. The Dewar was filled with liquid helium and an 8000 Gauss magnetic field applied. An extended series of experiments was carried out. Results were affected by the change in level of liquid helium in the Dewar. Addition of liquid nitrogen and refilling with liquid helium enormously increase the phonon level and affect the spin lattice relaxation time. The crystal does become opaque to red light on the time scale of the spin lattice relaxation time.

Conclusions

As a result of the extended series of measurements, it may be concluded that the collective absorption of red light by polarized nuclear spins is being observed. The selection rule derived in earlier reports--magnetic field vector of light parallel to applied magnetic field for greatest absorption--appears to be valid.

Starting with a one milliwatt polarized laser, the light is collimated, attenuated by neutral density filters to avoid heating the crystal, and permitted to interact with the crystal at low temperatures.

When the magnetic field is first applied immediately after cooldown, the light power transmitted to the photometer is 10^{-4} ergs per second. This drops by more than a factor ten, to a value too small to observe, on a time scale of approximately 24 hours. This confirms the earlier observations with an "unpolarized" laser.

6-86

1